

Identification and Characterization of *C. elegans* RNAi Inheritance Machinery

George Spracklin^{*,†}, Brandon Fields^{*,†}, Gang Wan[†], Diveena Vijayendran[†], Ashley Wallig^{*}, Aditi Shukla[†], and Scott Kennedy[†]

^{*} Laboratory of Genetics, University of Wisconsin-Madison, Madison, WI 53706

[†] Department of Genetics, Harvard Medical School, Boston, MA 02115

Data available from GEO under the accession number GSE92840

Running title: Identification and Characterization of *C. elegans* RNAi Inheritance Machinery

George Spracklin^{*†}, Brandon Fields^{*†}, Gang Wan[†], Diveena Vijayendran[†], Ashley Wallig^{*}, Aditi Shukla[†], and Scott Kennedy[†]

^{*} Laboratory of Genetics, University of Wisconsin-Madison, Madison, WI 53706

[†] Department of Genetics, Harvard Medical School, Boston, MA 02115

Keywords: epigenetic inheritance, small RNAs, *C.elegans*

Corresponding Author:

Scott Kennedy

77 Avenue Louis Pasteur

Boston, MA 02115

617-432-0094

kennedy@genetics.med.harvard.edu

ABSTRACT

Gene silencing mediated by dsRNA (RNAi) can persist for multiple generations in *Caenorhabditis elegans* (termed RNAi inheritance). Here we describe the results of a forward genetic screen in *C. elegans* that has identified six factors required for RNAi inheritance: GLH-1/VASA, PUP-1/CDE-1, MORC-1, SET-32, and two novel nematode specific factors that we term here (heritable RNAi defective) HRDE-2 and HRDE-4. The new RNAi inheritance factors exhibit mortal germline (Mrt) phenotypes, which we show is likely caused by epigenetic deregulation in germ cells. We also show that HRDE-2 contributes to RNAi inheritance by facilitating the binding of small RNAs to the inheritance Argonaute (Ago) HRDE-1. Together, our results identify additional components of the RNAi inheritance machinery whose conservation provides insights into the molecular mechanism of RNAi inheritance, further our understanding of how the RNAi inheritance machinery promotes germline immortality, and show that HRDE-2 couples the inheritance Ago HRDE-1 with the small RNAs it needs to direct RNAi inheritance and germline immortality.

INTRODUCTION

For most of the last century, it was widely believed that all epigenetic information was erased at, or near, the start of each new generation. It is now clear that this is not the case and that epigenetic information can be passed from parent to progeny. In some cases, epigenetic information can be transmitted from parent to progeny for multiple generations (termed transgenerational epigenetic inheritance, TEI). Examples of TEI include paramutation in plants, the inheritance of acquired traits in mammals, and RNA interference in nematodes (reviewed in Miska and Ferguson-Smith 2016). In many eukaryotes, dsRNA triggers silencing of cellular mRNAs, which contain sequence homology to the dsRNA, via a process known as RNA interference (RNAi). Interestingly, RNAi is heritable in *C. elegans*: the progeny of animals exposed to dsRNA continue to silence genes that were targeted by dsRNA in previous generations (termed RNAi inheritance). In some cases, RNAi inheritance can perdure for many generations (>5 generations) (Vastenhouw *et al.* 2006). RNAi inheritance is not accompanied by changes in DNA sequence, indicating that RNAi inheritance is an epigenetic phenomenon and a robust and unambiguous example of TEI.

dsRNAs are processed into small interfering RNAs (siRNAs) by the RNase III-like enzyme Dicer (Bernstein *et al.* 2001). siRNAs are bound by a conserved family of proteins termed the Argonautes (Ago). Together, Ago proteins and their associated siRNAs regulate gene expression by binding and inhibiting complementary cellular RNAs. In some organisms, like *C. elegans*, RNA-dependent RNA polymerases (RdRPs) use mRNA templates to amplify siRNA populations (termed 2° siRNAs) (Motamedi *et al.* 2004; Pak and Fire 2007; Sijen *et al.* 2007). In many animals, siRNAs enter the nucleus and regulate gene expression co-transcriptionally (termed nuclear RNAi) (Martienssen and Moazed 2015). In *C. elegans*, the nuclear-localized Argos HRDE-1 (germline) and NRDE-3 (soma) bind 2° siRNAs in the cytoplasm, move to the nucleus, and interact with complementary nascent transcripts via base pairing of their associated 2°

siRNAs. The nuclear Agos then recruit the nuclear RNAi factors (NRDE-1/2/4) to genomic sites of RNAi. Together, the nuclear RNAi machinery (by unknown mechanisms) directs the deposition of repressive chromatin marks such as H3K9me3 at RNAi-targeted loci and induce co-transcriptional inhibition of RNA Polymerase II (Guang *et al.* 2008; Guang *et al.* 2010; Burkhart *et al.* 2011). In the fission yeast *S. pombe*, nuclear RNAi triggers H3K9 methylation via the histone methyltransferase (HMT) Clr4 (Hall *et al.* 2002; Volpe *et al.* 2002). H3K9me contributes to nuclear RNAi by recruiting RNAi silencing factors to sites of nuclear RNAi (Motamedi *et al.* 2004). Thus, in *S. pombe*, non-coding RNAs and H3K9 methylation are part of a feed-forward loop that maintain stable states of nuclear gene silencing (Martienssen and Moazed 2015). It is not clear if or how RNAi-directed H3K9 methylation contributes to gene silencing in *C. elegans*.

Nuclear RNAi and RNAi inheritance require the same factors in *C. elegans* and nuclear gene-silencing events are needed for RNAi silencing signals to be inherited (Burton *et al.* 2011; Ashe *et al.* 2012; Buckley *et al.* 2012; Shirayama *et al.* 2012). Thus, nuclear RNAi and RNAi inheritance are related processes. HRDE-1 is an Ago that drives both nuclear RNAi and RNAi inheritance in germ cells (Buckley *et al.* 2012; Ashe *et al.* 2012; Shirayama *et al.* 2012; Luteijn *et al.* 2012). HRDE-1 is a member of an expanded clade of twelve *C. elegans* Agos (termed worm-specific Agos or WAGOs). The WAGOs are thought to bind endogenous small RNAs termed the 22G endo-siRNAs to mediate gene silencing of endogenous genes and RNAs (termed endogenous RNAi) (Billi *et al.* 2014). The degree to which the WAGOs act independently or redundantly during endogenous RNAi is unclear. It is also unclear if WAGOs bind the same siRNAs or if systems exist to direct subsets of 22G siRNAs to subsets of WAGOs.

HRDE-1 binds 22G endo-siRNAs that direct nuclear RNAi at hundreds of germline-expressed genes during the normal course of development (termed HRDE-1 target genes) (Buckley *et al.* 2012). HRDE-1 target genes include conserved genes, cryptic loci, and retrotransposons. Of these, LTR retrotransposons appear to be the most dramatically regulated loci (Ni *et al.* 2014). 21U RNAs, which are the *C. elegans* piRNAs, likely direct RdRPs to interact

with mRNAs to generate a subset of the 22G endo-siRNAs that bind HRDE-1 (Bagijn *et al.* 2012). However, most endo-siRNAs that associate with HRDE-1 are not piRNA dependent and the signals that direct RdRPs to produce these HRDE-1 siRNAs remain unknown (Ni *et al.* 2014). HRDE-1(-) animals (and animals lacking other known components of nuclear RNAi machinery) exhibit a mortal germline (Mrt) phenotype: Disabling the nuclear RNAi/RNAi inheritance machinery by mutation does not cause fertility defects immediately, however, after 3-5 generations mutant animals become sterile (Buckley *et al.* 2012). The following events occur coincidentally with the loss of germ cell viability, H3K9me3 marks are progressively lost at HRDE-1 target genes while expression of many HRDE-1 genes progressively increases (Buckley *et al.* 2012). Therefore, the RNAi inheritance machinery may promote germline immortality by maintaining H3K9me3 at HRDE-1 target genes to fine tune germline gene expression programs that are necessary for germ cell totipotency. The RNAi machinery is also known to silence parasitic genetic elements such as DNA transposons (Ketting *et al.* 1999). Therefore, it is also possible that Mrt could be caused by the accrual of genetic damage over generations as transposons or retrotransposons are mobilized in the germline of RNAi inheritance defective animals.

Here we report on the results of a genetic screen to identify factors required for RNAi inheritance in *C. elegans*. This screen identified six RNAi inheritance factors: GLH-1/VASA, a conserved RNA helicase, PUP-1/CDE-1, a nucleotidyltransferase, SET-32, a putative histone methyltransferase, MORC-1, a conserved GHKL ATPase, and two novel nematode specific factors that we term HRDE-2 and HRDE-4. We explore the role of these new RNAi inheritance factors in germline immortality and investigate in detail the mechanism by which one of these factors, HRDE-2, contributes to RNAi inheritance.

MATERIAL AND METHODS

Strains: N2, (YY513) *pkIs32[pie-1p::gfp::h2b]*, (YY528) *hrde-1(tm1200);pkIs32*, (YY1063) *hrde-2(gg517);pkIs32*, (YY673) *hrde-3(gg266);pkIs32*, (YY676) *hrde-4(gg300);pkIs32*, (YY494) *glh-1(ok439);pkIs32*, (YY1217) *pup-1(gg574);pkIs32*, (YY714) *morc-1(gg430);pkIs32;dpy-17(e164)*, (YY538) *hrde-1(tm1200)*, (YY1013) *hrde-2(gg517)*, (YY902) *set-32(gg266)*, (YY1197) *hrde-4(gg300)*, (YY976) *pup-1(gg519)*, (YY1195) *glh-1(gg408)*, (YY890) *morc-1(tm6048)*, (YY626) *hrde-2(gg268)*, (YY657) *hrde-1(tm1200);hrde-2(gg268)*, (YY909) *ggSi2[hrde-1p::3xflag::gfp::hrde-2];unc-119(ed3)*, (YY913) *nrde-2(gg518)*, (YY934) *hrde-2(gg517);nrde-2(gg518)*, (YY946) *rde-1(ne219);nrde-2(gg518)*, (YY947) *hrde-1(tm1200);nrde-2(gg518)*, (YY584) *ggSi1[hrde-1p::3xflag::gfp::hrde-1]*, (YY921) *rde-1(ne219);ggSi1*, (YY922) *hrde-2(gg517);ggSi1*, (YY636) *hrde-2(gg268);ggSi1*, (SX1316) *mjIs144[mex-5p::gfp::his-58::21UR-1::tbb-2 3'UTR];unc-119(ed3)*, (YY1041) *hrde-2(gg517);mjIs144*, (YY742) *hrde-2(gg252)*, (YY737) *hrde-2(gg399)*, (GR1747) *mut-15(tm1358)*, (YY626) *hrde-2(gg268);pkIs32*, (YY1143) *hrde-2(gg517);ggSi3[hrde-1p::gfp::hrde-2(AAAA)]*, (YY910) *hrde-2(gg252);ggSi2*, (YY634) *hrde-1(gg265);pkIs32*, (YY631) *hrde-1(gg257);pkIs32*, (YY562) *hrde-1(tm1200);oma-1(zu405)*, (TX20) *oma-1(zu405)*, (YY635) *hrde-2(gg268);oma-1(zu405)*, (SX461) *mjIs31[pie-1::gfp::h2b];unc-119(ed3)*, (YY720) *hrde-1(tm1200);mjIs31*, (YY722) *hrde-2(gg268);mjIs31*, (YY721) *hrde-1(tm1200);hrde-2(gg268);mjIs31*, (YY1246) *wago-3(gg589)*, (YY1340) *hrde-2(gg517);wago-3(gg589)*, (YY1312) *hrde-1(gg594)*, (YY1192) *hrde-1(gg594);hrde-2(gg517)*

Construction of plasmids and transgenic strains: For 3XFLAG::GFP::HRDE-1 (*ggSi1*) (referred to as FLAG::HRDE-1 when assaying HRDE-1 immunoprecipitation) the *hrde-1* promoter, coding region, and predicted 3' UTR were amplified by PCR from genomic N2 DNA and cloned into pCFJ151 using Gibson cloning. For GFP::3XFLAG::HRDE-2 (*ggSi2*), the *hrde-2*

coding region and predicted 3'UTR, along with the *hrde-1* promoter were amplified by PCR and assembled using Gibson cloning into pCFJ151. DEVD sequence (A.A. 99-102) in HRDE-2 was mutated to AAAA in *ggSi3*. The forementioned plasmids were injected into EG6699 and integrated into the genome using the MosSCI technique (Frøkjær-Jensen et al. 2008). Insertions were verified using PCR and sequencing.

NRDE-2::3XFLAG::HA (*nrde-2(gg518)*) (referred to as FLAG::NRDE-2 when assaying NRDE-2 immunoprecipitations) CRISPR strain was generated with 50ng/ul pDD162 (Cas9), 25ng/ul pRB1017-derived guide RNA, and 500nM single-stranded oligo (IDT 4-nmol Ultramer) using protocol as previously described (Farboud and Meyer 2015). A gRNA targeting *unc-58* and repair oligo (AF-JA-76) were used as a co-conversion marker. Insertions were verified using PCR and sequencing.

CRISPR deletions (*hrde-2(gg517)*, *set-32(gg545)*, *set-32(gg546)*, *hrde-4(gg558)*, *pup-1(gg519)*) were generated using the protocol above with the exception of an extra gRNA were added to cause two dsDNA breaks.

RNAi: RNAi experiments were conducted as previously described (Timmons and Fire 1998; Burton *et al.* 2011). All clones were taken from the Ahringer library (Kamath *et al.* 2003).

RNA IP: Adult worms were flash frozen then sonicated (2 min, 30s ON 30s OFF 30% amplitude) three times in sonication buffer (20mM Tris pH 7.5, 200mM NaCl, 2.5mM MgCl₂, 0.5% NP-40, 10% glycerol, RNaseOUT, Proteinase Inhibitor(Roche)) in a QSonica 800R sonicator. Samples were then rotated for 15 min at 4°C then spun at 14k, 15min, 4°C. Lysates were pre-cleared with 30ul Protein-A/Salmon Sperm DNA agarose beads (16-157, Millipore) for 30min at 4°C then spun 14k, 5 min, 4°C. Supernatant was then transferred to fresh tube with 2ul of anti-flag M2 conjugated to agarose (A2220, Sigma) and rotated at 4°C for 4hrs-O/N. Six washes with 1mL RIP buffer [20mM Tris pH 7.5, 200mM NaCl, 2.5mM MgCl₂, 0.5% NP-40, 10% glycerol] spin at 2K, 1min.

After washes, 100ul of elution buffer (1mL sonication buffer, 3ul 3xFLAG peptide (5 mg/mL, F4799 Sigma)) was then added and rotated 30min at 4°C. Supernatant transferred to new tube, 5ul saved for protein assays, the sample was then DNaseI treated (add MgCl₂), 37°C, 20min. Samples were then added to 400ul TRIzol and standard isopropanol precipitation. Resuspended in 20ul dH₂O.

Chromatin IP: Gravid adult animals were cross-linked in 2% formaldehyde/M9 solution for 30 min at room temperature, and fixation was stopped by adding glycine to 0.125M and incubating for 5 min. Gravid adult animals were washed twice with M9 and stored in -80°C. Fixed gravid adult animals were resuspended in 1 ml FA buffer (50 mM Tris/HCl pH 7.5, 1 mM EDTA, 1% Triton X-100, 0.1% sodium deoxycholate, 150 mM NaCl, with proteinase inhibitor (Roche)). Samples were sonicated 30s ON 30s OFF 70% amplitude for 20 mins in a QSonica 800R sonicator. Lysates were clarified by centrifugation for 10 minutes at 4°C. Supernatant was transferred to a new tube and pre-cleared with salmon sperm DNA-coated protein-A agarose beads (16-157, Millipore). Pre-cleared lysates were incubated with 2 µl anti- trimethylated Histone 3 lysine 9 (Upstate 07-523) for 2hrs - O/N. BSA and ssDNA pre-treated protein-A agarose beads were added. Beads were pelleted and washed twice in FA buffer, twice in FA-500 buffer (50 mM Tris/HCl pH 7.5, 1 mM EDTA, 1% Triton X-100, 0.1% sodium deoxycholate, 500 mM NaCl), once in LiCl wash buffer (0.25 M LiCl, 1% NP-40, 1% sodium deoxycholate, 1 mM EDTA, 10 mM Tris-HCl, pH 8.0), and once in TE buffer. Chromatin was eluted with elution buffer (1% SDS, 0.1M NaHCO₃) for 30 minutes each at room temperature. 30 ul 5 M NaCl was added, and reversed crosslinks by heating at 65°C O/N. DNAs were further purified by QIAquick PCR purification kit (Qiagen), and analyzed by quantitative real-time PCR (qPCR). ChIP data was normalized to co-immunoprecipitated *eft-3*.

cDNA Preparation: RNAs were converted to cDNA using the iScript cDNA Synthesis Kit (Bio-Rad, 170-8890) following manufacturer's protocol.

TaqMan: Total RNA was isolated from adult worms by vortexing flash frozen extracts for 10min in TRIzol, followed by isopropanol precipitation. *oma-1* small RNAs were reverse-transcribed by microRNA reverse-transcription kits (4366596; Applied Biosystems). cDNA was quantified by qPCR using custom Taqman small RNA assay from Applied Biosystems (4398987; assay IDs CSKAJ9W, CSLJIF4, CSMSGMC, CSN1ESK). Taqman sequences were the same as used in (Buckley *et al.* 2012).

Small RNA cloning and Computational Analysis: Small RNAs were enriched with a mirVana miRNA Isolation Kit (Ambion, AM1560). 2ug of small RNAs were cloned using a 5'-mono-phosphate independent small RNA protocol as previously described by (Gent *et al.* 2009). Libraries were multiplexed with a 4nt 5' barcode and a 6nt 3' barcode and pooled for HTS on a NextSeq500 (Biopolymers Facility, HMS).

Fastx 0.0.13 was used to separate reads that contained the 3' adapter and filter low quality reads for further analysis. Custom scripts were used to separate samples based on the in-line 5' barcode. Reads >17nt were mapped to the *C. elegans* genome (WS220) using Bowtie 1.1.1 '-f -v 3 -p 4 -a --best --strata -m 400' (described in (Shirayama *et al.* 2014)). The read counts were normalized to the total number of reads that matched the genome (in millions). To account for differences in sequencing volume between samples, reads were normalized too non-structural RNAs. Custom scripts were used to draw the single nucleotide plots. All scripts available upon request.

Accession numbers: Illumina data are available from GEO under the accession number GSE92840.

Mrt assay: At each generation, 6 larval stage (L3) animals were picked to a single plate. Average brood sizes were calculated by counting the total number of progeny per plate.

***oma-1* RNAi assay:** 6 larval stage (L1) were singled onto bacteria expressing *oma-1* dsRNA and propagated at 20°C (restrictive temperature). These animals were designated the P0/RNAi generation. 6 L1 progeny (F1) animals were then transferred to bacteria without *oma-1* dsRNA. Brood size was obtained from counting F1 gravid adults remaining on the P0 plates. This process was repeated at every generation.

Primers: *oma-1* 1F/R: (TTGTTAAGCATTCCCTGCAC) and (TCGATCTTCTCGTTGTTTTCA)
oma-1 2F/R: (GTGAGTTTAACAAAATTCAAATGAAAA) and (AGTGGTGAAACGGGCAAAGT)
spr-2 RT 1F/R: (AACTTTTCGCAATGTA ACTCTTCC) and (CTGGTAGTGCTGCGTACGTG)
C09G9.5 RT 1F/R: (CCAACGAAGCTTTGTGGATA) and (GCGTGTGGGAACCTAGAAAA)
C27B7.2 1F/R: (TGGGTAACTGAAACA ACTTGAA) and (TCACAAACAACGTCTTTTCGAT)
crs-2: (GCCGGAAGTTGAAA ACTACATC) and (TTCGTCCAAATCCTCGAATC)
Y61A9LA.12: TGGAGAGCACTCTGAAA AATTG) and (TCTATTCCAGCAATTTCTTATCTCAA)
bath-45: (CTGATTCAAAGCTATTCGGAGAA) and (CATCGATCAGGAAAGCATCA)
R11A8.1: (TCTTCGGCGCTAATCTTTTC) and (TTGGTAGAAGCTGCATCACTTT)
tlk-1: (TCATTTTGGCGACTTCATCA) and (TTTGTTATTGAATGTGACGTG)
lit-1: (TCAATTTTTGCCATGGAATC) and (ATGTCCTTCTCCCACTTTGG)
eft-3: (ACTTGATCTACAAGTGCGGAGGA) and (CGGGTGAGAAAATCTTTCAA ACTA)

RESULTS

A genetic screen identifies RNAi inheritance factors: Exposure of wild-type *C. elegans* to *gfp* dsRNA causes silencing of a germline-expressed *gfp* transgene and this silencing is heritable (Figure 1) (Vastenhouw *et al.* 2006; Buckley *et al.* 2012). To further our understanding of the mechanism of RNAi inheritance, we conducted a forward genetic screen looking for mutations that specifically disrupted the inheritance phase of RNAi. Briefly, animals expressing GFP in germ cells were EMS mutagenized. F2 progeny of mutagenized animals were exposed to *gfp* dsRNA. We isolated mutant animals that were able to silence GFP in response to *gfp* RNAi but failed to inherit GFP silencing in subsequent generations. We screened ~6,000,000 haploid genomes and identified 57 heritable RNAi defective (*hrde*) alleles (Figure S1). Previously, this screen identified the nuclear Ago HRDE-1/WAGO-9 (henceforth referred to as HRDE-1) and the nuclear RNAi factors NRDE-1/2/4 (Buckley *et al.* 2012). The following approaches were used to focus our efforts on novel RNAi inheritance factors. First, animals lacking NRDE-1/2/4 respond idiosyncratically to dsRNAs targeting the *pos-1* and *lin-26* genes (Burkhart *et al.* 2011; Buckley *et al.* 2012). Mutant strains that behaved like *nrde-1/2/4* mutant animals, when treated with *pos-1* and *lin-26* RNAi, were discarded as many of these animals likely harbored mutations in *nrde-1/2/4*. Second, *hrde* alleles were complemented with *hrde-1(tm1200)* animals, and those alleles that failed to complement *hrde-1* were not analyzed further (Figure S2). The remaining *hrde* alleles (23/57) were subjected to whole genome sequencing (WGS). Combinations of the following approaches, which varied for each *hrde* gene, were used to assign molecular identities: positional mapping, complementation analysis, genetic rescue, and *in silico* complementation analysis (Minevich *et al.* 2012) (see below). Finally, additional mutant alleles (generated by CRISPR/CAS9, or obtained from the Caenorhabditis Genome Center) of candidate *hrde* genes were tested for RNAi inheritance defects in order to confirm identities. Evidence establishing the identity of each *hrde* gene is summarized in Table 1 and is described in detail in Figure S2-S8. In addition, a detailed

description of the identification of *hrde-2* is presented in the main text below. Together, these experiments show that GLH-1/VASA, PUP-1/CDE-1, SET-32, MORC-1, and two nematode specific factors HRDE-2 and HRDE-4, are required for RNAi inheritance in *C. elegans* (Figure 1). The causative mutations in three *hrde* strains remain to be identified.

RNAi inheritance factors promote germline immortality: The germline is immortal in the sense that, unlike somatic cells, germ cells and their descendents are able to divide in perpetuity. Germline immortality is under genetic control (Ahmed and Hodgkin 2000; Buckley *et al.* 2012; Simon *et al.* 2014; Sakaguchi *et al.* 2014). Interestingly, the previously known components of the RNAi inheritance machinery are required for germline immortality: animals that lack NRDE-1/2/4 or HRDE-1 show near wild-type fertility, however, the descendents of these animals become sterile after 3-5 generations at 25°C (Buckley *et al.* 2012). This transgenerational loss of fertility is referred to as a mortal germline (Mrt) phenotype (Ahmed and Hodgkin 2000). We asked if animals lacking the newly identified components of the RNAi inheritance machinery were Mrt. Note, the Mrt phenotype of RNAi inheritance defective animals (such as *nrde-2* or *hrde-1* mutants) is temperature sensitive: growth at 25°C induces Mrt, however, growth at 20°C does not induce Mrt (Buckley *et al.* 2012). Therefore, we shifted *glh-1*, *pup-1*, *set-32*, *morc-1*, *hrde-2*, and *hrde-4* mutant animals to growth at 25°C and monitored the fertility of these animals over generations. As expected, wild-type animals were not Mrt, and *hrde-1* mutant animals were Mrt (Figure 2). *glh-1*, *pup-1*, *set-32*, *morc-1*, *hrde-2*, and *hrde-4* mutant animals were Mrt (Figure 2). We tested a second allele of each of these genes (except for *morc-1*) and found that these strains also exhibited Mrt phenotypes. These data show that a general property of RNAi inheritance factors is to promote germline immortality.

The RNAi inheritance machinery likely promotes germline immortality by maintaining the *C. elegans* epigenome: The RNAi machinery regulates the expression of both protein-coding

genes and parasitic nucleic acids such as transposons (Meister 2013). Therefore, the Mrt phenotype of RNAi inheritance defective animals might be caused by progressive changes in gene expression or by genetic damage resulting from transposon mobilization. The following experiments show that the cause of germline mortality in RNAi inheritance defective animals is unlikely to be genetic damage. After 2-3 generations of growth at the non-permissive (25°C) temperature, the brood size of *hrde-1* or *hrde-2* mutant animals was <25% that of wild-type animals (Figure 3A). We transferred these low-fertility animals to the permissive temperature (20°C) and found that after three generations, fertility in these populations was restored to near wild-type levels (Figure 3A and Figure S9). Moving these animals back to the non-permissive temperature again resulted in a progressive decline in fertility, which, again, could be reversed by a shift to permissive temperatures (Figure 3A and Figure S9). These data show that the Mrt phenotype of *hrde-1* or *hrde-2* mutant animals is reversible and these data are consistent with recently published reports (Simon *et al.* 2014; Sakaguchi *et al.* 2014; Ni *et al.* 2016). We considered the possibility that rare individuals, which had not incurred DNA damage, might be present in *hrde* populations undergoing Mrt. These animals might, over generations, overtake sicker DNA-damaged siblings, thus, reversing Mrt. To address this issue, we quantified the brood of 84 *hrde-1/-2* double mutant animals from populations that were almost Mrt at permissive temperatures. We found that animals from a Mrt population exhibited a spectrum of fertility rates ranging from zero to near wild-type (Figure 3B, top). Importantly, individuals in the population that had the lowest (non-zero) brood sizes gave rise to progeny that had wild-type fertility (Figure 3B, bottom). We conclude that Mrt is reversible and that the accumulation of DNA damage does not cause Mrt in RNAi inheritance defective animals.

How then does the RNAi inheritance machinery promote germline immortality? It has been proposed that endogenously expressed siRNAs may promote germline immortality by maintaining a germline H3K9me3 pattern that is necessary for germ cell health (Buckley *et al.* 2012). Our inheritance screen identified nine alleles of the gene *set-32*, which encodes a SET domain protein

with homology to *S. pombe* Clr4 and fly Su(var)3-9. Clr4 and Su(var)3-9 are founding members of a family of enzymes that methylate H3K9 (Rea *et al.* 2000). Thus, SET-32 is a candidate H3K9 methyltransferase. Additionally, *set-32* mutant animals are Mrt, consistent with the idea that maintenance of germline H3K9me3 patterns by SET-32 is required for germline immortality (Figure 2). HRDE-1 is the Ago that engages nuclear RNAi to maintain germline immortality and the gene targets of HRDE-1 are known (Buckley *et al.* 2012). To test the idea that SET-32-directed H3K9me3 may promote germline immortality, we asked if SET-32 contributed to H3K9me3 at HRDE-1 target genes. Animals lacking SET-32 had less H3K9me3 at HRDE-1 target genes than wild-type animals (Figure 4A). H3K9me3 at control (non-HRDE-1 target) genes was unaffected by loss of SET-32 (Figure 4A). Exposing *C. elegans* to experimental dsRNAs mimics the effects of endogenous siRNAs and induces H3K9me3 at genomic sites that exhibit sequence homology to dsRNAs (Guang *et al.* 2010). SET-32 was required for experimentally supplied *oma-1* dsRNA to trigger H3K9me3 at the *oma-1* locus (Figure 4B). Thus, SET-32 is required for H3K9me3 to accumulate at genes targeted by exogenous and endogenous sources of siRNAs in *C. elegans*. These data suggest that SET-32 is an siRNA-directed H3K9 methyltransferase and that siRNA-directed H3K9 methylation is an obligate step during RNAi inheritance. Additionally, these data support the model that the RNAi inheritance machinery promotes germline immortality by establishing an epigenome conducive to germ cell function.

Molecular identification of *hrde-2*: The remainder of this paper describes the identification and characterization of *hrde-2*. *hrde-2* was identified in our genetic screen for heritable silencing of a *gfp* transgene. We wondered if HRDE-2 was required for RNAi inheritance at endogenous genes. *oma-1* and *pos-1* are two germline expressed genes and RNAi silencing of these genes is heritable. *hrde-2* animals were defective for the inheritance of *oma-1* and *pos-1* RNAi gene silencing, indicating that HRDE-2 is generally required for RNAi inheritance in *C. elegans* (Figure S10A,B). The following lines of evidence show that *hrde-2* is *t01c3.9*, a nematode specific factor

required for RNAi inheritance. We mapped *hrde-2* to a 2 cM interval on chromosome V (Figure S3B). Within this interval, we identified loss of function mutations in the open reading frame of *t01c3.9* in three out of our three *hrde-2* strains, which previous work had established were members of the *hrde-2* complementation group (Figure 5A, Table 1, and Figure S3A). We generated a deletion in *t01c3.9* with CRISPR/Cas9 and found that animals harboring this deletion allele were defective for RNAi inheritance (Figure 1). Finally, a wild-type copy of *t01c3.9* was sufficient to rescue the Mrt phenotype of *hrde-2* mutant animals (Figure S11). We conclude that *hrde-2* is *t01c3.9*, a nematode specific protein that is required for RNAi inheritance. Note, *t01c3.9* is co-transcribed in an operon with the gene *mut-15/rde-5*, which is known to be involved in RNAi silencing in *C. elegans* (Sijen *et al.* 2007; Phillips *et al.* 2012). *hrde-2(gg268)* complemented *mut-15(tm1358)* for RNAi defects, indicating that mutation of *t01c3.9* does not cause RNAi inheritance defects by affecting expression of *mut-15* (Figure S3C).

HRDE-2 localizes to cytoplasmic foci in germ cells: The following observations argue that HRDE-2 is a germline-expressed RNAi inheritance factor. First, we created a *gfp::hrde-2* fusion gene, which we integrated into the genome using MosSCI technology (Frøkjær-Jensen *et al.* 2008). *gfp::hrde-2* rescued the Mrt phenotype of *hrde-2* mutant animals, suggesting that *gfp::hrde-2* encodes a functional protein whose expression likely reflects that of endogenous HRDE-2 (Figure S11). We observed GFP::HRDE-2 fluorescence throughout the male and female germline in *gfp::hrde-2* animals (Figure 5B and Figure S12). GFP::HRDE-2 appeared concentrated in perinuclear foci that surrounded nuclei (see discussion). In addition, microarray and RNA-seq based studies show that the *hrde-2* mRNA expression is enriched/restricted to the germline (Reinke *et al.* 2004; Ortiz *et al.* 2014). Together, these data argue that HRDE-2 is a germline-expressed RNAi inheritance factor.

HRDE-2 is a component of the *C. elegans* nuclear RNAi pathway in the germline: The *C. elegans* nuclear RNAi machinery is required for RNAi inheritance (Burton *et al.* 2011; Buckley *et al.* 2012; Ashe *et al.* 2012; Shirayama *et al.* 2012). We wondered if HRDE-2 might function as part of the nuclear RNAi pathway. To address this question, we first conducted a genetic analysis with *hrde-2* and the nuclear RNAi Ago *hrde-1* to ask if these genes acted in the same genetic pathway. Animals lacking both HRDE-1 and HRDE-2 became Mrt at a similar rate as animals lacking either HRDE-1 or HRDE-2, individually (Figure 6A). This data suggests that *hrde-2* acts in the same genetic pathway as *hrde-1* to promote germline immortality.

The following data provide further evidence that HRDE-2 is a component of nuclear RNAi pathway. First, piRNA-mediated gene silencing of a *gfp* reporter transgene (*mjls144*) requires nuclear RNAi (Bagijn *et al.* 2012). We found that, like the known components of the nuclear RNAi pathway, HRDE-2 was required for piRNA-mediated gene silencing of this reporter gene (Figure 6B). Second, as described above, exposing *C. elegans* to dsRNAs is sufficient to induce H3K9me3 at genomic sites that exhibit sequence homology to dsRNAs (Guang *et al.* 2010). We found that, like the other known components of the nuclear RNAi pathway, HRDE-2 was required for *oma-1* dsRNA to trigger H3K9 methylation at the *oma-1* locus (Figure 6C and Figure S13). The nuclear RNAi factor NRDE-2 binds nascent RNAs (pre-mRNA) when genes are targeted by dsRNA, and this step of nuclear RNAi requires the Ago HRDE-1 (Guang *et al.* 2010; Buckley *et al.* 2012). We found that, like HRDE-1, HRDE-2 was required for dsRNA to induce NRDE-2/pre-mRNA association (Figure 6D). Similarly, we found that HRDE-2 was required for NRDE-2 to bind the pre-mRNA of an endogenous target of nuclear RNAi, *bath-45* (Figure 6E). Taken together, these data show that HRDE-2 is a component of the nuclear RNAi pathway and that HRDE-2 acts upstream of NRDE-2/pre-mRNA binding and NRDE-dependent H3K9 methylation during nuclear RNAi.

HRDE-2 is required for the Ago HRDE-1 to bind siRNAs following RNAi: HRDE-2 acts in the nuclear RNAi pathway upstream of NRDE-2/pre-mRNA binding and NRDE-dependent H3K9 methylation. HRDE-1 is the major Ago that drives germline nuclear RNAi in *C. elegans* (Buckley *et al.* 2012). To further pinpoint how HRDE-2 contributes to nuclear RNAi, we asked if HRDE-2 was required for HRDE-1 to bind siRNAs. HRDE-1 binds endogenous small RNAs (termed 22G siRNAs), which are produced by RdRPs acting upon cytoplasmic mRNA templates (Pak and Fire 2007; Sijen *et al.* 2007; Buckley *et al.* 2012). For unknown reasons, 22G siRNAs are predominantly antisense to the RNAs used by RdRPs for their production. We treated wild-type and *hrde-2* mutant animals with *oma-1* dsRNA, immunoprecipitated HRDE-1, and isolated co-precipitating RNAs. We then used four sets of TaqMan probes, which were designed to detect *oma-1* 22G siRNAs (antisense to the *oma-1* mRNA), to ask if HRDE-2 was required for HRDE-1 to associate with siRNAs. Two of these Taqman primer sets detect antisense *oma-1* siRNAs derived from sequences covered by the *oma-1* dsRNA and two primer sets detect antisense *oma-1* siRNAs derived from sequences not covered by the *oma-1* dsRNA. Using all four probe sets, we found that HRDE-1 failed to associate with *oma-1* siRNAs in *hrde-2* mutant animals (Figure 7A and Figure S14). Note: Expression of the HRDE-1 was reduced (1.4-fold) in a *hrde-2* mutant background (Figure S15), however, this reduction is unlikely to explain the 60-fold reduction in HRDE-1 co-precipitating small RNAs that we observe in *hrde-2* mutant animals (Figure 7A). Together, these data suggest that the reason HRDE-2 is required for RNAi inheritance is that HRDE-2 is required for the Ago HRDE-1 to bind siRNAs.

HRDE-1 is one of twelve members of a clade of *C. elegans* Agos collectively referred to as the worm-specific Agos or WAGOs (Yigit *et al.* 2006). Like HRDE-1, the other eleven WAGOs are thought to bind RdRP-produced 22G siRNAs (Yigit *et al.* 2006; Gu *et al.* 2009). We wondered if HRDE-2 was generally required for all WAGOs to bind siRNAs or if HRDE-2 function might be restricted to a subset of the WAGOs. To address this question, we used CRISPR/Cas9 to tag the endogenous *wago-3/ppw-2* locus (hereafter referred to as WAGO-3) with an HA-epitope and then

used this reagent to ask if HRDE-2 was required for WAGO-3 to associate with siRNAs. We chose WAGO-3 for these experiments because, like HRDE-1, WAGO-3 is expressed in the germline (data not shown). After exposure to *oma-1* dsRNA, HA::WAGO-3 associated with similar amounts of *oma-1* siRNAs in wild-type and *hrde-2* mutant animals (Figure 7A). These data show that the role of HRDE-2 in WAGO/siRNA binding is limited to a subset of the WAGOs.

HRDE-2 might be required for producing HRDE-1 siRNAs or for facilitating the binding of HRDE-1 to siRNAs. To begin to address this question, we sequenced total small RNAs from wild-type and *hrde-2* mutant animals that were exposed to *oma-1* dsRNA or empty vector. Wild-type and *hrde-2* mutant animals produced similar levels of *oma-1* siRNAs (Figure 7B). The similarity in *oma-1* siRNA patterns extended to regions of the *oma-1* mRNA not directly targeted by *oma-1* dsRNA, establishing that RdRP activity is present and apparently normal in *hrde-2* mutant animals (Figure 7B). In empty vector treated animals, very few *oma-1* siRNAs were identified (<0.5% of *oma-1* RNAi treated animals) (Figure S16). These results indicate that *hrde-2* mutant animals do not exhibit obvious defects in generating the type of siRNA to which HRDE-1 binds. Taken together, our results suggest that the function of HRDE-2 is to facilitate the binding of HRDE-1 to the siRNA cofactors it needs to direct RNAi inheritance and germline immortality.

DISCUSSION

Here we report the identification of six RNAi inheritance factors; GLH-1/VASA, CDE-1/PUP-1, SET-32, MORC-1, HRDE-2, and HRDE-4. HRDE-2 is required for the Ago HRDE-1 to bind siRNAs after RNAi treatment. Interestingly, HRDE-2 does not appear to be needed for the production of the siRNAs to which HRDE-1 binds, suggesting that the function of HRDE-2 is to assist HRDE-1 in binding siRNAs. It is unclear if HRDE-2 has a direct or indirect role in promoting this HRDE-1/siRNA interaction. In *S. pombe*, the Argonaute small interfering RNA chaperone (ARC) complex facilitates Ago1 siRNA binding (Holoch and Moazed 2015). *C. elegans* HRDE-2 may function analogously to ARC complex to directly mediate the loading of siRNAs into HRDE-1. Alternatively, HRDE-2 may promote HRDE-1/siRNA binding by an indirect mechanism. For instance, HRDE-2 could be needed to activate a factor that itself is required for HRDE-1/siRNA binding. Favoring (but not proving) this model, is the fact that at steady state, HRDE-2 localizes to the cytoplasm (Figure 5B and Figure S12) while HRDE-1 localizes to the nucleus (Buckley *et al.* 2012; Shirayama *et al.* 2012). Future *in vivo* interaction and *in vitro* biochemical studies may help differentiate these models. Interestingly, we find that HRDE-2 is not required for other Agos (WAGO-3) to bind siRNAs in response to dsRNA treatment, indicating some degree of specificity for HRDE-2 function during Ago/siRNA binding. *C. elegans* possess ~27 Agos and more than twelve of these Agos (WAGOs) are thought to bind chemically indistinguishable 22G siRNAs. If or how WAGOs associate with subsets of the total cellular pool of 22G siRNAs is unknown. Our results suggest that HRDE-2 could be one mechanism by which different WAGOs are coupled to different 22G siRNAs. Interestingly, the *C. elegans* genome encodes two paralogs of *hrde-2* (*t24c2.2* and *f43c11.9*), which may act as specificity factors for other *C. elegans* Argonaute proteins.

H3K9me3, RNAi inheritance, and germline immortality: There are 38 potential histone-tail methyltransferases (HMTs) encoded in the *C. elegans* genome (Andersen and Horvitz 2007). Our genetic screen identified nine alleles of one of these HMTs, SET-32. These results suggest that SET-32 is the major HMT contributing to germline RNAi inheritance in *C. elegans*. SET-32 is required for H3K9me3 to accumulate at genomic sites targeted by endogenous and exogenous siRNAs (Figure 4). Together, these data argue that SET-32 is likely an H3K9 methyltransferase and that siRNA-directed H3K9 methylation is likely an important step for transgenerational gene silencing. In addition, we show that SET-32 is required for germline immortality, suggesting that siRNA-directed H3K9 methylation is likely an important facet of small RNA-mediated germline immortality.

How might SET-32 promote RNAi inheritance? In *S. pombe*, Ago1 directs the HMT Clr4 to genomic sites targeted by siRNAs so that Clr4 can methylate H3K9 (Volpe *et al.* 2002). Additionally, the chromodomain protein Chp1 binds H3K9me3 and recruits Ago1 to genomic sites of RNAi (Partridge *et al.* 2002; Verdell *et al.* 2004; Hayashi *et al.* 2012; Rougemaille *et al.* 2012). Thus, in *S. pombe*, small RNAs and H3K9 methylation act together in a feed-forward loop to maintain stable states of nuclear gene silencing (Martienssen and Moazed 2015). It is possible that a similar feed-forward mechanism may be at play in *C. elegans*. The nuclear RNAi machinery could direct SET-32 to chromatin to deposit H3K9me3, which could then help stabilize the nuclear RNAi machinery at repressive chromatin. It will, therefore, be of interest to ask if SET-32 physically associates with chromatin in response to RNAi, and if SET-32 and the nuclear RNAi factors require each other for stable association with chromatin. Finally, it may be worth noting that our screen identified a mutation in SET-32 that mutates a serine residue to phenylalanine (Ser170Phe), hinting that SET-32 might be regulated by phosphorylation. If this turns out to be the case, such a system might provide an elegant mechanism by which *C. elegans* could fine tune RNAi inheritance programs in response to changing environmental conditions.

VASA, germ granules, and RNAi inheritance: VASA is a conserved DEAD-box RNA helicase that contributes to germ cell function in most metazoans (Voronina *et al.* 2011). Here we show that *C. elegans* lacking GLH-1/VASA are defective for RNAi inheritance (Figure 1). In insects, VASA helps to silence transposable elements (TEs) by binding TE RNAs and processing TE RNAs into intermediates that are fed to Ago proteins (Guzzardo *et al.* 2013; Xiol *et al.* 2014). The association between VASA-like helicases and Ago proteins may be conserved: RDE-12, Belle, and MVH are all VASA homologues that associate with Ago proteins in worms, flies, and mice, respectively (Kirino *et al.* 2010; Pek and Kai 2011; Shirayama *et al.* 2014; Yang *et al.* 2014). Together, these observations hint that GLH-1/VASA helicases may be conserved mediators of small RNA-based gene silencing pathway in metazoans and that VASA-like enzymes might fulfill this role by linking RNAi factors to RNAs in need of silencing.

A previous study failed to detect RNAi defects in *glh-1/vasa* mutant animals (Spike *et al.* 2008). We speculate that this is likely due to the fact that, while *glh-1/vasa* mutant animals are defective for RNAi inheritance, they respond normally when exposed directly to dsRNA (Figure 1). Why would the loss of GLH-1/VASA specifically affect the inheritance phase of RNAi? Germ granules are self-assembling non-membrane bound organelles found in the germ cells of most metazoans (Voronina *et al.* 2011). In *C. elegans*, germ granules are referred to as P granules. P granules segregate with the germ lineage during embryonic cell divisions (Strome and Wood 1983). GLH-1 (like VASA homologues in most animals) localizes to germ granules (Voronina *et al.* 2011). The concentration of GLH-1/VASA into P granules, and the segregation of these granules with the germline, position GLH-1/VASA perfectly to mediate heritable gene silencing events. It is important to note, however, that GLH-1/VASA activity is required for normal P granule formation (Spike *et al.* 2008). Several core RNAi factors (such as Dicer, Agos, and RdRPs) localize to P granules (Gu *et al.* 2009; Claycomb *et al.* 2009; Conine *et al.* 2010; Vasale *et al.* 2010; Beshore *et al.* 2011). It is possible, therefore, that failure to properly concentrate core RNAi factors in P granules could be responsible for the RNAi inheritance defects observed in GLH-1(-)

animals. Consistent with this latter idea, DEPS-1 and PGL-1 are two germline factors that are required for wild-type P granule assembly and animals lacking either of these factors exhibit RNAi defects (Spike *et al.* 2008; Robert *et al.* 2005).

RNA uridylation and RNAi inheritance: Poly(u) polymerase-1 (*pup-1*, also known as *cde-1* or *cid-1*) encodes a poly(U) polymerase that adds short tracts of uracil to the 3' termini of RNAs in *C. elegans* (Kwak and Wickens 2007; van Wolfswinkel *et al.* 2009). PUP-1 is thought to uridylate the endo-siRNAs that bind a germline Ago called CSR-1 (van Wolfswinkel *et al.* 2009). PUP-1 uridylation is thought to destabilize CSR-1 siRNAs (van Wolfswinkel *et al.* 2009). Thus, in animals lacking PUP-1, more CSR-1 siRNAs are available for binding to CSR-1. Unlike most Agos (which negatively regulate target RNAs), CSR-1, and its associated siRNAs, are thought to activate gene expression (Cecere *et al.* 2014; Wedeles *et al.* 2013; Seth *et al.* 2013). Therefore, one model to explain why PUP-1 is required for RNAi inheritance is the following: in *pup-1* mutant animals, CSR-1 is hyperactive and the enhanced anti-silencing activity of CSR-1 prevents RNAi inheritance by overriding the pro-silencing activity of the RNAi inheritance machinery. Alternatively, PUP-1 may also play a direct role in RNAi inheritance. In *S. pombe*, a homologue of PUP-1 (Cid12) is part of a complex of proteins termed the RNA directed RNA polymerase complex (RDRC). RDRC consists of Cid12, as well as an RNA helicase (Hrr1), and an RNA dependent RNA Polymerase (RdRP) (Motamedi *et al.* 2004). The RDRC is thought to produce silencing RNAs needed to perpetuate stable long-term heterochromatic states in *S. pombe* (Martienssen and Moazed 2015). Thus, PUP-1-like enzymes have now been associated with long-term gene silencing processes in both yeast and worms. Perhaps RNA uridylation in *C. elegans* is a signal that shunts RNAs into long-term gene silencing pathways. For instance, in somatic cells of *C. elegans*, the putative poly(U) polymerase RDE-3 is thought to add a uracil to the 3' end of mRNA fragments produced in response to RNAi (Tsai *et al.* 2015). It has been suggested that these 3' uracils may serve as beacons to recruit RdRPs, which amplify small RNA

populations and reinforce gene silencing (Tsai *et al.* 2015). PUP-1 could function analogously by marking germline RNAs as templates for RdRPs activity each generation during RNAi inheritance. By so doing, PUP-1 could maintain small RNA populations at levels sufficient to trigger gene silencing over multiple generations.

MORC-1: MORC-1 is a member of a conserved family of GHKL ATPases named after the mouse Microorchidia (*Morc*) gene. In plants, MORC family members help silence repetitive genomic elements such as TEs. Plant MORCs are thought to be recruited to TE genes by RNA-directed DNA methylation (RdDM) (Liu *et al.* 2014). Once localized to TEs, MORC enzymes may regulate gene expression by compacting chromatin into higher order structures that are not conducive to transcription (Moissiard *et al.* 2012). Interestingly, in mice, MORC also repress TEs, hinting that the silencing of genome parasites may be conserved feature of the MORC family of ATPases (Pastor *et al.* 2014). The *C. elegans* MORC homologue (MORC-1) has been previously implicated in RNAi silencing processes (Robert *et al.* 2005; Kim *et al.* 2005; Moissiard *et al.* 2012). Our results suggest that MORC-1 activity may be particularly important during the inheritance phase of RNAi. If MORC-1 were a component of the nuclear RNAi machinery one would expect such an inheritance specific defect. Perhaps nuclear RNAi directs MORC-1 to genes undergoing heritable gene silencing in *C. elegans* in order to compact chromatin into states not compatible with gene expression. Asking if nuclear RNAi compacts chromatin in *C. elegans*, and if MORC-1 is needed for this compaction, will be an important test of this model.

REFERENCES

- Ahmed S., Hodgkin J., 2000 MRT-2 checkpoint protein is required for germline immortality and telomere replication in *C. elegans*. *Nature* 403: 159–164.
- Alcazar R. M., Lin R., Fire A. Z., 2008 Transmission dynamics of heritable silencing induced by double-stranded RNA in *Caenorhabditis elegans*. *Genetics* 180: 1275–1288.
- Andersen E. C., Horvitz H. R., 2007 Two *C. elegans* histone methyltransferases repress *lin-3* EGF transcription to inhibit vulval development. *Development* 134: 2991–2999.
- Ashe A., Sapetschnig A., Weick E.-M., Mitchell J., Bagijn M. P., *et al.*, 2012 piRNAs can trigger a multigenerational epigenetic memory in the germline of *C. elegans*. *Cell* 150: 88–99.
- Bagijn M. P., Goldstein L. D., Sapetschnig A., Weick E.-M., Bouasker S., *et al.*, 2012 Function, targets, and evolution of *Caenorhabditis elegans* piRNAs. *Science* 337: 574–578.
- Bernstein E., Caudy A. A., Hammond S. M., Hannon G. J., 2001 Role for a bidentate ribonuclease in the initiation step of RNA interference. *Nature* 409: 363–366.
- Beshore E. L., McEwen T. J., Jud M. C., Marshall J. K., Schisa J. A., *et al.*, 2011 *C. elegans* Dicer interacts with the P-granule component GLH-1 and both regulate germline RNPs. *Dev. Biol.* 350: 370–381.
- Billi A. C., Fischer S. E. J., Kim J. K., 2014 Endogenous RNAi pathways in *C. elegans*. *WormBook*: 1–49.
- Buckley B. A., Burkhart K. B., Gu S. G., Spracklin G., Kershner A., *et al.*, 2012 A nuclear Argonaute promotes multigenerational epigenetic inheritance and germline immortality. *Nature* 489: 447–451.

- Burkhart K. B., Guang S., Buckley B. A., Wong L., Bochner A. F., *et al.*, 2011 A pre-mRNA-associating factor links endogenous siRNAs to chromatin regulation. *PLoS Genet.* 7: e1002249.
- Burton N. O., Burkhart K. B., Kennedy S., 2011 Nuclear RNAi maintains heritable gene silencing in *Caenorhabditis elegans*. *Proc. Natl. Acad. Sci. U. S. A.* 108: 19683–19688.
- Cecere G., Hoersch S., O’Keeffe S., Sachidanandam R., Grishok A., 2014 Global effects of the CSR-1 RNA interference pathway on the transcriptional landscape. *Nat. Struct. Mol. Biol.* 21: 358–365.
- Claycomb J. M., Batista P. J., Pang K. M., Gu W., Vasale J. J., *et al.*, 2009 The Argonaute CSR-1 and its 22G-RNA cofactors are required for holocentric chromosome segregation. *Cell* 139: 123–134.
- Conine C. C., Batista P. J., Gu W., Claycomb J. M., Chaves D. A., *et al.*, 2010 Argonautes ALG-3 and ALG-4 are required for spermatogenesis-specific 26G-RNAs and thermotolerant sperm in *Caenorhabditis elegans*. *Proc. Natl. Acad. Sci. U. S. A.* 107: 3588–3593.
- Farboud B., Meyer B. J., 2015 Dramatic enhancement of genome editing by CRISPR/Cas9 through improved guide RNA design. *Genetics* 199: 959–971.
- Frøkjær-Jensen C., Davis M. W., Hopkins C. E., Newman B. J., Thummel J. M., *et al.*, 2008 Single-copy insertion of transgenes in *Caenorhabditis elegans*. *Nat. Genet.* 40: 1375–1383.
- Gent J. I., Schvarzstein M., Villeneuve A. M., Gu S. G., Jantsch V., *et al.*, 2009 A *Caenorhabditis elegans* RNA-directed RNA polymerase in sperm development and endogenous RNA interference. *Genetics* 183: 1297–1314.

- Gu W., Shirayama M., Conte D. Jr, Vasale J., Batista P. J., *et al.*, 2009 Distinct argonaute-mediated 22G-RNA pathways direct genome surveillance in the *C. elegans* germline. *Mol. Cell* 36: 231–244.
- Guang S., Bochner A. F., Pavelec D. M., Burkhart K. B., Harding S., *et al.*, 2008 An Argonaute transports siRNAs from the cytoplasm to the nucleus. *Science* 321: 537–541.
- Guang S., Bochner A. F., Burkhart K. B., Burton N., Pavelec D. M., *et al.*, 2010 Small regulatory RNAs inhibit RNA polymerase II during the elongation phase of transcription. *Nature* 465: 1097–1101.
- Guzzardo P. M., Muerdter F., Hannon G. J., 2013 The piRNA pathway in flies: highlights and future directions. *Curr. Opin. Genet. Dev.* 23: 44–52.
- Hall I. M., Shankaranarayana G. D., Noma K.-I., Ayoub N., Cohen A., *et al.*, 2002 Establishment and maintenance of a heterochromatin domain. *Science* 297: 2232–2237.
- Hayashi A., Ishida M., Kawaguchi R., Urano T., Murakami Y., *et al.*, 2012 Heterochromatin protein 1 homologue Swi6 acts in concert with Ers1 to regulate RNAi-directed heterochromatin assembly. *Proc. Natl. Acad. Sci. U. S. A.* 109: 6159–6164.
- Holoch D., Moazed D., 2015 Small-RNA loading licenses Argonaute for assembly into a transcriptional silencing complex. *Nat. Struct. Mol. Biol.* 22: 328–335.
- Kamath R. S., Fraser A. G., Dong Y., Poulin G., Durbin R., *et al.*, 2003 Systematic functional analysis of the *Caenorhabditis elegans* genome using RNAi. *Nature* 421: 231–237.
- Ketting R. F., Haverkamp T. H., Luenen H. G. van, Plasterk R. H., 1999 Mut-7 of *C. elegans*, required for transposon silencing and RNA interference, is a homolog of Werner syndrome helicase and RNaseD. *Cell* 99: 133–141.

- Kim J. K., Gabel H. W., Kamath R. S., Tewari M., Pasquinelli A., *et al.*, 2005 Functional genomic analysis of RNA interference in *C. elegans*. *Science* 308: 1164–1167.
- Kirino Y., Vourekas A., Kim N., Lima Alves F. de, Rappsilber J., *et al.*, 2010 Arginine methylation of vasa protein is conserved across phyla. *J. Biol. Chem.* 285: 8148–8154.
- Kwak J. E., Wickens M., 2007 A family of poly(U) polymerases. *RNA* 13: 860–867.
- Lin R., 2003 A gain-of-function mutation in *oma-1*, a *C. elegans* gene required for oocyte maturation, results in delayed degradation of maternal proteins and embryonic lethality. *Dev. Biol.* 258: 226–239.
- Liu Z.-W., Shao C.-R., Zhang C.-J., Zhou J.-X., Zhang S.-W., *et al.*, 2014 The SET domain proteins SUVH2 and SUVH9 are required for Pol V occupancy at RNA-directed DNA methylation loci. *PLoS Genet.* 10: e1003948.
- Luteijn M. J., Bergeijk P. van, Kaaij L. J. T., Almeida M. V., Roovers E. F., *et al.*, 2012 Extremely stable Piwi-induced gene silencing in *Caenorhabditis elegans*. *EMBO J.* 31: 3422–3430.
- Martienssen R., Moazed D., 2015 RNAi and heterochromatin assembly. *Cold Spring Harb. Perspect. Biol.* 7: a019323.
- Meister G., 2013 Argonaute proteins: functional insights and emerging roles. *Nat. Rev. Genet.* 14: 447–459.
- Minevich G., Park D. S., Blankenberg D., Poole R. J., Hobert O., 2012 CloudMap: a cloud-based pipeline for analysis of mutant genome sequences. *Genetics* 192: 1249–1269.
- Miska E. A., Ferguson-Smith A. C., 2016 Transgenerational inheritance: Models and mechanisms of non-DNA sequence-based inheritance. *Science* 354: 59–63.

- Moissiard G., Cokus S. J., Cary J., Feng S., Billi A. C., *et al.*, 2012 MORC family ATPases required for heterochromatin condensation and gene silencing. *Science* 336: 1448–1451.
- Motamedi M. R., Verdell A., Colmenares S. U., Gerber S. A., Gygi S. P., *et al.*, 2004 Two RNAi complexes, RITS and RDCR, physically interact and localize to noncoding centromeric RNAs. *Cell* 119: 789–802.
- Ni J. Z., Chen E., Gu S. G., 2014 Complex coding of endogenous siRNA, transcriptional silencing and H3K9 methylation on native targets of germline nuclear RNAi in *C. elegans*. *BMC Genomics* 15: 1157.
- Ni J. Z., Kalinava N., Chen E., Huang A., Trinh T., *et al.*, 2016 A transgenerational role of the germline nuclear RNAi pathway in repressing heat stress-induced transcriptional activation in *C. elegans*. *Epigenetics Chromatin* 9: 3.
- Ortiz M. A., Noble D., Sorokin E. P., Kimble J., 2014 A new dataset of spermatogenic vs. oogenic transcriptomes in the nematode *Caenorhabditis elegans*. *G3* 4: 1765–1772.
- Pak J., Fire A., 2007 Distinct populations of primary and secondary effectors during RNAi in *C. elegans*. *Science* 315: 241–244.
- Partridge J. F., Scott K. S. C., Bannister A. J., Kouzarides T., Allshire R. C., 2002 cis-acting DNA from fission yeast centromeres mediates histone H3 methylation and recruitment of silencing factors and cohesin to an ectopic site. *Curr. Biol.* 12: 1652–1660.
- Pastor W. A., Stroud H., Nee K., Liu W., Pezic D., *et al.*, 2014 MORC1 represses transposable elements in the mouse male germline. *Nat. Commun.* 5: 5795.
- Pek J. W., Kai T., 2011 DEAD-box RNA helicase Belle/DDX3 and the RNA interference pathway promote mitotic chromosome segregation. *Proc. Natl. Acad. Sci. U. S. A.* 108: 12007–12012.

- Phillips C. M., Montgomery T. A., Breen P. C., Ruvkun G., 2012 MUT-16 promotes formation of perinuclear mutator foci required for RNA silencing in the *C. elegans* germline. *Genes Dev.* 26: 1433–1444.
- Rea S., Eisenhaber F., O'Carroll D., Strahl B. D., Sun Z. W., *et al.*, 2000 Regulation of chromatin structure by site-specific histone H3 methyltransferases. *Nature* 406: 593–599.
- Reinke V., Gil I. S., Ward S., Kazmer K., 2004 Genome-wide germline-enriched and sex-biased expression profiles in *Caenorhabditis elegans*. *Development* 131: 311–323.
- Robert V. J. P., Sijen T., Wolfswinkel J. van, Plasterk R. H. A., 2005 Chromatin and RNAi factors protect the *C. elegans* germline against repetitive sequences. *Genes Dev.* 19: 782–787.
- Rougemaille M., Braun S., Coyle S., Dumesic P. A., Garcia J. F., *et al.*, 2012 Ers1 links HP1 to RNAi. *Proc. Natl. Acad. Sci. U. S. A.* 109: 11258–11263.
- Sakaguchi A., Sarkies P., Simon M., Doebley A.-L., Goldstein L. D., *et al.*, 2014 *Caenorhabditis elegans* RSD-2 and RSD-6 promote germ cell immortality by maintaining small interfering RNA populations. *Proc. Natl. Acad. Sci. U. S. A.* 111: E4323–31.
- Seth M., Shirayama M., Gu W., Ishidate T., Conte D. Jr, *et al.*, 2013 The *C. elegans* CSR-1 argonaute pathway counteracts epigenetic silencing to promote germline gene expression. *Dev. Cell* 27: 656–663.
- Shirayama M., Seth M., Lee H.-C., Gu W., Ishidate T., *et al.*, 2012 piRNAs initiate an epigenetic memory of nonself RNA in the *C. elegans* germline. *Cell* 150: 65–77.
- Shirayama M., Stanney W. 3rd, Gu W., Seth M., Mello C. C., 2014a The Vasa Homolog RDE-12 engages target mRNA and multiple argonaute proteins to promote RNAi in *C. elegans*. *Curr. Biol.* 24: 845–851.

- Sijen T., Steiner F. A., Thijssen K. L., Plasterk R. H. A., 2007 Secondary siRNAs result from unprimed RNA synthesis and form a distinct class. *Science* 315: 244–247.
- Simon M., Sarkies P., Ikegami K., Doebley A.-L., Goldstein L. D., *et al.*, 2014 Reduced insulin/IGF-1 signaling restores germ cell immortality to *Caenorhabditis elegans* Piwi mutants. *Cell Rep.* 7: 762–773.
- Spike C. A., Bader J., Reinke V., Strome S., 2008a DEPS-1 promotes P-granule assembly and RNA interference in *C. elegans* germ cells. *Development* 135: 983–993.
- Spike C., Meyer N., Racen E., Orsborn A., Kirchner J., *et al.*, 2008b Genetic analysis of the *Caenorhabditis elegans* GLH family of P-granule proteins. *Genetics* 178: 1973–1987.
- Strome S., Wood W. B., 1983 Generation of asymmetry and segregation of germ-line granules in early *C. elegans* embryos. *Cell* 35: 15–25.
- Timmons L., Fire A., 1998 Specific interference by ingested dsRNA. *Nature* 395: 854.
- Tsai H.-Y., Chen C.-C. G., Conte D. Jr, Moresco J. J., Chaves D. A., *et al.*, 2015 A ribonuclease coordinates siRNA amplification and mRNA cleavage during RNAi. *Cell* 160: 407–419.
- Vasale J. J., Gu W., Thivierge C., Batista P. J., Claycomb J. M., *et al.*, 2010 Sequential rounds of RNA-dependent RNA transcription drive endogenous small-RNA biogenesis in the ERGO-1/Argonaute pathway. *Proc. Natl. Acad. Sci. U. S. A.* 107: 3582–3587.
- Vastenhouw N. L., Brunschwig K., Okihara K. L., Müller F., Tijsterman M., *et al.*, 2006 Gene expression: long-term gene silencing by RNAi. *Nature* 442: 882.
- Verdel A., Jia S., Gerber S., Sugiyama T., Gygi S., *et al.*, 2004 RNAi-mediated targeting of heterochromatin by the RITS complex. *Science* 303: 672–676.

- Volpe T. A., Kidner C., Hall I. M., Teng G., Grewal S. I. S., *et al.*, 2002 Regulation of heterochromatic silencing and histone H3 lysine-9 methylation by RNAi. *Science* 297: 1833–1837.
- Voronina E., Seydoux G., Sassone-Corsi P., Nagamori I., 2011 RNA granules in germ cells. *Cold Spring Harb. Perspect. Biol.* 3.
- Wedeles C. J., Wu M. Z., Claycomb J. M., 2013 Protection of germline gene expression by the *C. elegans* Argonaute CSR-1. *Dev. Cell* 27: 664–671.
- Wolfswinkel J. C. van, Claycomb J. M., Batista P. J., Mello C. C., Berezikov E., *et al.*, 2009 CDE-1 affects chromosome segregation through uridylation of CSR-1-bound siRNAs. *Cell* 139: 135–148.
- Xiol J., Spinelli P., Laussmann M. A., Homolka D., Yang Z., *et al.*, 2014 RNA clamping by Vasa assembles a piRNA amplifier complex on transposon transcripts. *Cell* 157: 1698–1711.
- Yang H., Vallandingham J., Shiu P., Li H., Hunter C. P., *et al.*, 2014 The DEAD box helicase RDE-12 promotes amplification of RNAi in cytoplasmic foci in *C. elegans*. *Curr. Biol.* 24: 832–838.
- Yigit E., Batista P. J., Bei Y., Pang K. M., Chen C.-C. G., *et al.*, 2006 Analysis of the *C. elegans* Argonaute family reveals that distinct Argonautes act sequentially during RNAi. *Cell* 127: 747–757.

Figure Legends

Figure 1. Genetic screen identifies *hrde* genes. Animals of the indicated genotype expressing a *pie-1p::gfp::h2b* transgene were exposed to bacteria producing *gfp* dsRNA. F1 embryos were isolated by hypochlorite treatment and grown on bacteria not expressing *gfp* dsRNA. GFP expression in oocytes of animals exposed to *gfp* dsRNA (RNAi generation) and the progeny of these animals (inheriting generation) were visualized by fluorescence microscopy. Percentage of animals expressing GFP is indicated (n>50). Note: *morc-1* is marked with *dpy-17(e164)*.

Figure 2. *hrde* genes are required for germline immortality. Brood sizes for animals of indicated genotype were counted as detailed in material and methods. Data are mean \pm s.d. (n=3).

Figure 3. Genetic damage is unlikely the cause of Mrt in RNAi inheritance defective animals. (A) Brood sizes for animals of indicated genotype were counted as detailed in material and methods. Shift in temperature is indicated by changing color scheme (n=6, \pm s.d.). **(B) *hrde-1;hrde-2* animals from the 5th generation at 25°C were singled and grown at 20°C and their brood sizes were scored (B, top).** Animals whose parents had a low brood size (indicated by red bars) were singled, grown at 20°C, and their brood sizes were scored **(B, bottom)**.

Figure 4. SET-32 is required for nuclear RNAi-directed H3K9me3. (A) H3K9me3 ChIP was conducted on animals of indicated genotype as described in materials and methods. qPCR of H3K9me3 co-precipitating DNA using primers detecting 5 genes thought to be targeted by nuclear RNAi and 3 genes not thought to be targeted by nuclear RNAi. Data are relative to co-precipitating *eft-3* DNA and expressed as a ratio of signal from indicated mutant to signal from wild-type (n=6, \pm s.d.). **(B)** Animals exposed to *oma-1* dsRNA (+ *oma-1* dsRNA) or control bacteria with L4440 vector (- *oma-1* dsRNA) were subjected to H3K9me3 ChIP. Co-precipitating *oma-1* DNA was

quantified by qPCR. Data were normalized to co-precipitating *eft-3* DNA and expressed as a ratio of signals in animals subjected to *oma-1* RNAi over signals from no RNAi (-) animals ('1' denotes no change) (n=3, \pm s.d). On the x-axis, 0 denotes the predicted start codon of *oma-1*.

Figure 5. *hrde-2* encodes a germline factor required for RNAi inheritance. (A) Predicted *hrde-2* gene structure. Arrows indicate mutant alleles identified in genetic screen and black bar indicates deletion allele created by CRISPR/CAS9. **(b)** HRDE-2 localizes to cytoplasmic foci in germ cells. Fluorescence microscopy of the hermaphrodite germline in adult animals expressing *gfp::hrde-2*. Arrow shows perinuclear puncta.

Figure 6. HRDE-2 is part of the nuclear RNAi pathway. (A) Brood sizes for animals of indicated genotype were counted as detailed in material and methods. *hrde-1;hrde-2* double mutant animals do not show an additive fertility defect at 25°C when compared to the single mutant animals, suggesting HRDE-1 and HRDE-2 act in the same pathway. (n=6, \pm s.d) **(B)** Fluorescence images of oocytes from wild-type animals expressing a piRNA sensor transgene (*mjls144*), which contains a binding site for the piRNA 21U-1, and a control piRNA sensor transgene (*mjls145*) which lacks the 21U-1 site (as described in *Bagijn et al., 2012*). Right panel, *hrde-2(gg517)* animals harboring *mjls145* (top) and *mjls144* (bottom). **(C)** Animals of the indicated genotypes were exposed to *oma-1* dsRNA (+ *oma-1* dsRNA) or empty L4440 vector (- *oma-1* dsRNA) and subjected to H3K9me3 ChIP. Co-precipitating *oma-1* DNA was quantified by qPCR. Data were normalized to co-precipitating *eft-3* DNA and expressed as a ratio \pm *oma-1* RNAi ('1' denotes no change) (n=3, \pm s.d). On the x-axis, 0 denotes the predicted start codon of *oma-1*. **(D/E)** FLAG-NRDE-2 was precipitated from animals of indicated genotypes with an anti-Flag antibody and NRDE-2 co-precipitating RNA was isolated by Trizol extraction. RNA was converted to cDNA and quantified by qRT-PCR with primers detected *oma-1* pre-mRNA or *bath-45* pre-mRNA as described in materials and methods. Comparisons were performed using *t*-tests (Prism) **(E)** Data

are expressed as fold change \pm *oma-1* RNAi, or **(E)** as fold change mutant/control. Mean \pm s.d. are shown (n=3.).

Figure 7. HRDE-2 is required for HRDE-1 to associate with siRNAs. **(A)** FLAG::HRDE-1 or HA::WAGO-3 expressing animals were treated with *oma-1* dsRNA (+*oma-1* dsRNA) or with empty L4440 vector (-*oma-1* dsRNA). HRDE-1 and WAGO-3 were immunoprecipitated with anti-Flag and anti-HA antibodies respectively, and co-precipitating RNA was isolated by TRIzol extraction. *oma-1* co-precipitating siRNAs were quantified with an *oma-1* TaqMan probe set that detected a 22G RNA antisense to the *oma-1* mRNA. TaqMan signals shown relative to Taqman signals from non-RNAi treated control animals, which was defined as one. Relative enrichment following RNAi treatment is shown (n=3, \pm s.d.). Comparisons were performed using *t*-tests (Prism) **(B)** Wild-type and *hrde-2(gg517)* animals were treated with *oma-1* dsRNA. Small RNAs were sequenced as described in materials and methods. Small RNAs that mapped to the *oma-1* locus are shown. Sense siRNA (red) and antisense siRNA (blue). Data were normalized by the total number of sequenced reads for each genotype, reads per million (rpm). *oma-1* siRNAs sequenced from control (non RNAi) samples are shown in Figure S16. A cartoon representing the *oma-1* genomic regions is shown below the graphs. The region of the *oma-1* locus targeted by dsRNA is demarcated with a red line and a black box.

Figure S1. Genetic screen for hrde alleles. *pie-1p::gfp::h2b (pkIs32)* expressing animals were mutagenized with EMS. F2 progeny of these mutagenized animals were subjected to *gfp* RNAi. F3 progeny were scored for GFP expression. Mutant animals unable to inherit *gfp* RNAi silencing were kept for further analysis. As a secondary screen, we tested mutant animals for their ability to silence *gfp*, *pos-1*, and *lin-26* dsRNA. We discarded mutants that failed to silence *gfp* or *pos-1* when exposed directly to dsRNA (likely representing animals with global defects in RNAi silencing (*rde*)). Next we used *lin-26* RNAi in an attempt to identify and eliminate mutants harboring alleles

of *nrde-1/2/4*. *lin-26* RNAi induces a lethal phenotype in wild-type animals. *nrde-1/2/3/4* mutant animals fail to respond to *lin-26* RNAi when directly exposed to *lin-26* dsRNA. Mutant animals that were defective for *lin-26* RNAi were discarded. Note: it is possible that these secondary screens may have prevented our identification of some genes involved in multi-generational RNAi inheritance. Of the remaining 122 mutant strains, we chose a single mutant from each pool, which exhibited highly (>90%) penetrant RNAi inheritance defects (57 total) for further analysis.

¹Additional screens used *dpy-17(e164);pkIs32* and *dpy-17(e164);pkIs32;neSi21[3xflag:hrde-1]*

as the starting strain. ²Some additional screens put F3 animals onto RNAi opposed to F2 animals.

(Adapted from Buckley *et al.*, 2012)

Figure S2. *hrde-1* is required for RNAi inheritance *hrde-1* was the first gene identified by this genetic screen and reported in Buckley *et al.*, 2012 **(a)** Predicted domain structure of HRDE-1 protein is shown. *hrde-1* mutant alleles (missense only) are indicated by arrows. **(b)** List of identified mutations for each allele which are grouped based on missense, nonsense, or unidentified. Sanger sequencing identified mutations within the *c16c10.3* coding regions in 19 strains. Complementation analyses support the identification of 6 additional alleles. N/A, Not Available.

Figure S3. *t01c3.9* is required for RNAi inheritance **(a)** List of alleles and identified mutations within the *t01c3.9* coding region. Supporting evidence for each allele (RNAi inheritance defects, H3K9me3 defects after exposure to *oma-1* dsRNA, and Mrt phenotype) is reported. ND, not done. **(b)** *hrde-2* mutant animals expressing *gfp::h2b* were crossed with a mapping strain (*gfp::h2b* transgene backcrossed to CB4856 (Hawaiian)). Recombinant F2 animals were exposed to *gfp* dsRNA and F3 animals that were defective for GFP silencing were isolated and genotyped. We mapped *hrde-2* to a 2 cM interval on LG V. **(c)** *hrde-2(gg268)* and *mut-15(tm1358)* mutant animals

expressing *gfp::h2b* were mated and heterozygous progeny were exposed to *gfp* dsRNA. GFP was visualized by fluorescent microscopy.

Figure S4. *set-32* is required for RNAi inheritance (a) Predicted *set-32* gene structure. Arrows indicate mutant alleles. Black bar indicates deletion allele. (b) List of alleles and identified mutations within the *set-32* coding region. Supporting evidence for each allele (RNAi inheritance defects, H3K9me3 defects after exposure to *oma-1* dsRNA, and Mrt phenotype) is reported. ND, not done.

Figure S5. *t10b11.7* is required for RNAi inheritance (a) Predicted *hrde-4* gene structure. Arrows indicate mutant alleles. Black bar indicates deletion allele. (b) List of alleles and identified mutations within the *t10b11.7* coding region. Supporting evidence for each allele (RNAi inheritance defects, H3K9me3 defects after exposure to *oma-1* dsRNA, and Mrt phenotype) is reported. ND, not done.

Figure S6. *glh-1* is required for RNAi inheritance (a) Predicted *glh-1* gene structure. Arrow indicates mutant allele. Black bar indicates deletion allele. (b) List of alleles and identified mutations within the *glh-1* coding region. Supporting evidence for each allele (RNAi inheritance defects, H3K9me3 defects after exposure to *oma-1* dsRNA, and Mrt phenotype) is reported. ND, not done.

Figure S7. *pup-1* is required for RNAi inheritance (a) Predicted *pup-1* gene structure. Arrows indicate mutant alleles. Black bars indicate deletion alleles. (b) List of alleles and identified mutations within the *pup-1* coding region. Supporting evidence for each allele (RNAi inheritance defects, H3K9me3 defects after exposure to *oma-1* dsRNA, and Mrt phenotype) is reported. ND, not done.

Figure S8. *morc-1* is required for RNAi inheritance (a) Predicted *morc-1* gene structure. Arrows indicate mutant alleles. Black bar indicates deletion allele. (b) List of alleles and identified mutations within the *morc-1* coding region. Supporting evidence for each allele is reported (RNAi inheritance defects, H3K9me3 defects after exposure to *oma-1* dsRNA, and Mrt phenotype). ND, not done.

Figure S09. Non-genetic damage likely driving progressive sterility in *hrde* mutant animals. Brood sizes of the indicated genotypes were scored across generations. Temperature shift are denoted in blue=20°C or pink=25°C. Data are mean \pm s.d. (n=6). Middle and Bottom graphs are continuations of the graphs above them.

Figure S10. *hrde-2* is required for RNAi inheritance (a) Animals of the indicated genotypes were exposed to *pos-1* dsRNA at a concentration that produced >50% F₁ lethality. Surviving F₁ progeny were grown in the absence of *pos-1* dsRNA and % F₂ embryonic arrest was scored. (b) *zu405ts* is a temperature sensitive (ts) lethal (embryonic arrest at 20°C) allele of *oma-1* (Lin 2003). *oma-1* RNAi suppresses *oma-1(zu405ts)* lethality, and this effect is heritable (Alcazar, Lin, and Fire 2008). P₀ animals were exposed to *oma-1* RNAi and number of viable progeny in the indicated generations were scored (20°C) (n=3, \pm s.d.). Note: *hrde-2(gg268)* animals are partially defective for inheritance of *oma-1* silencing suggesting that HRDE-2 helps promote RNAi inheritance, albeit weaker than HRDE-1.

Figure S11. GFP::*T01C3.9* rescues defects in *hrde-2* mutant animals An exogenous copy of *hrde-2* is sufficient of rescuing the brood size defects associated with *hrde-2* mutant animals at elevated temperatures, confirming the molecular identity of *hrde-2*. *t01c3.9* fused to *gfp* (*gfp::hrde-2*) was integrated as a single copy transgene into the genome using MosSCI. *gfp::hrde-2* was

crossed into *hrde-2(gg252)* and *hrde-2(gg517)* backgrounds then shifted to the restricted temperature to test for rescue of the Mrt phenotype.

Figure S12. *gfp::hrde-2* is expressed during spermatogenesis and in the male germline GFP::*HRDE-2* was visualized using fluorescent microscopy in the male germline. Arrow shows perinuclear puncta.

Figure S13. HRDE-2 is required for RNAi-directed H3K9me3 Animals of the indicated genotypes were exposed to *oma-1* dsRNA (+*oma-1* dsRNA) or with empty vector (-*oma-1* dsRNA) and subjected to H3K9me3 ChIP. Co-precipitating *oma-1* DNA was quantified by qPCR. Data were normalized to co-precipitating *eft-3* DNA and expressed as a ratio \pm *oma-1* RNAi ('1' denotes no change) (n=3, \pm s.d). On the x-axis, primer pairs were located within the transcriptional unit (*oma-1* and *spr-2*), or outside the transcriptional unit (*rad-26*).

Figure S14. HRDE-2 is required for HRDE-1 to associate with siRNAs (a/b/c) FLAG::*HRDE-1* or HA::*WAGO-3* expressing animals were treated with *oma-1* dsRNA (+*oma-1* dsRNA) or with empty vector (-*oma-1* dsRNA). *HRDE-1* and *WAGO-3* were immunoprecipitated with anti-Flag and anti-HA antibodies, respectively. Co-precipitating RNA was isolated by TRIzol extraction. *oma-1* co-precipitating siRNAs were quantified with an *oma-1* TaqMan probe set that detected a 22G RNA antisense to the *oma-1* mRNA. TaqMan signals shown relative to Taqman signals from non-RNAi treated control animals, which was defined as one. Relative enrichment following RNAi treatment is shown (n=3, \pm s.d.) **(d)** Schematic of *oma-1* locus with TaqMan probes labeled A-D and region targeted by dsRNA labelled in red. Data for probe A is in Figure 7.

Figure S15. GFP::*HRDE-1* is not mislocalized in *hrde-2* mutant animals nor is expression of *HRDE-1* dramatically affected by loss of *HRDE-2*. **(a)** *HRDE-1* is localized to germ cell nuclei

(Ashe et al. 2012; Buckley et al. 2012). 3XFLAG::GFP::HRDE-1 (*ggSi1*) localizes to germ cell nuclei and is not mislocalized in *hrde-2(gg268)* mutant animals. GFP::HRDE-2 was visualized using fluorescent microscopy. Arrows indicate nuclei. **(b/c)** Western blot detecting HA::WAGO-3 and FLAG::HRDE-1 and, as a loading control, tubulin. **(b)** YY1246 (lane 1) and YY1340 (lane 2) **(c)** YY584 (lane 1), YY922 (lane 2), YY636 (lane 3), and YY921 (lane 4) animals were boiled in Laemmli buffer and samples were subjected to SDS PAGE electrophoresis followed by probing with **(b)** α -HA or **(c)** α -FLAG and **(b/c)** α -tubulin antibodies. Band intensities were quantified with ImageJ software.

Figure S16. HRDE-2 is not required for small RNA biogenesis. Wild-type and *hrde-2(gg517)* animals were treated with *oma-1* dsRNA. Small RNAs were sequenced as described in materials and methods. Small RNAs that mapped to the *oma-1* locus are shown. Sense siRNA (red) and antisense siRNA (blue). Data were normalized by the total number of sequenced reads for each genotype reads per million (rpm). *oma-1* siRNAs sequenced from control (non RNAi) samples. A cartoon representing the *oma-1* genomic regions is shown below the graphs. The region of the *oma-1* locus targeted by dsRNA is demarcated with a red line and a black box. *oma-1* dsRNA treated sample are from Figure 7.

Table 1: Genetic screen identifies seven genes as required for RNAi inheritance. ^a Number of alleles identified from the forward genetic screen outlined in Figure S1. ^b Independently isolated alleles in the corresponding gene (obtained via CGC or CRISPR/Cas9). ^c Defining members of each complementation group were determined by exposing F1 trans-heterozygotes to *gfp* dsRNA and assaying percentage of F2 animals that failed to inherit GFP silencing.

	Gene	Predicted Function	Alleles Identified ^a	Additional Alleles ^b	Complementation Group ^c
<i>hrde-1</i>	<i>wago-9</i>	Argonaute	25	<i>tm1200</i>	<i>tm1200, gg265, gg257</i>
<i>hrde-2</i>	<i>t01c3.9</i>	Unknown	4	<i>gg517</i>	<i>gg268, gg252, gg399</i>
<i>hrde-3</i>	<i>set-32</i>	H3K9 methyltransferase	9	<i>gg545, gg546, ok1457</i>	<i>gg374, gg375, gg376</i>
<i>hrde-4</i>	<i>t10b11.7</i>	Unknown	4	<i>gg558</i>	<i>in silico</i>
<i>hrde-5</i>	<i>glh-1</i>	VASA/Helicase	1	<i>ok439</i>	N/A
<i>hrde-6</i>	<i>pup-1</i>	poly(U)polymerase	1	<i>tm936, tm1021, gg519</i>	N/A
<i>hrde-7</i>	<i>morc-1</i>	ATPase	1	<i>tm6048, gk1754232</i>	N/A




Figure 1





Figure 2

A



B




Figure 3






Figure 4

A



B




Figure 5





Figure 6

A



B




Figure 7